

欧文原著論文

1. Kawamura T, Egusa G, Fujikawa R, and Okubo M.
 β_3 -adrenergic receptor gene variant is associated with upper body obesity only in obese Japanese-American men but not in women.
Diabetes Res Clin Pract. 54:49-55. 2001.
2. Kawamura T, Egusa G, Fujikawa R, and Okubo M.
Gln27Glu variant of the β_2 -adrenergic receptor gene is not associated with Obesity and Diabetes in Japanese-Americans.
Metabolism. 50:443-446. 2001.
3. Kawamura T, Egusa G, Okubo M, Imazu M, Yamakido M.
Association of β_3 -adrenergic receptor gene polymorphism with insulin resistance in Japanese-American men.
Metabolism. 48:1367-70. 1999.
4. Kawamura T, Egusa G, Fujikawa R, Watanabe T, Oda K, Kataoka S, Takayama S, Kubo K, Yamamoto S, Noma K, Orita R, and Yamakido M.
Effect of acarbose on glycemic control and lipid metabolism in patients with non-insulin-dependent diabetes mellitus.
Current therapeutic research. 59 : 97-106, 1998.
5. Okubo M, Watanabe H, Fujikawa R, Kawamura T, Egusa G, Yamakido M.
Reduced prevalence of diabetes according to 1997 American Diabetes Association criteria.
Diabetologia. 42:1168-70. 1999.

欧文症例報告論文

1. Kawamura T, Nimura I, Hanafusa M, Fujikawa R, Okubo M, Egusa G, Yamakido M.
DiGeorge Syndrome with Graves' Disease: A Case Report.
Endocrine Journal. 47:91-95, 2000.
2. Yamashita Y, Kawamura T, Fujikawa R, Mochizuki H, Okubo M, Arita K.
Regression of both pituitary and ovarian cysts after administration of thyroid hormone in a case of primary hypothyroidism.
Intern Med. 40:751-5, 2001.